HIPP COMBIOTIK®

Preparation











Follow the instructions very carefully. Incorrect preparation and storage of the feed can promote growth of unwanted bacteria.

- 1. Boil fresh water from tap to 100°C for a few minutes. Cool down to 40-50° C.
- 2. Add the correct number of scoops of powder for one meal into a clean bottle.
- 3. Add water to the bottle. Place the cap on the bottle and shake well.
- 4. Cool down to drinking temperature. Test the temperature of the milk on the inside of your wrist before feeding.
- 5. Serve immediately and discard any leftovers after the meal

Dosing chart

| Boiled water | Level scoops | Prepared feed |
|--------------|--------------|---------------|
| 60 ml | 2 | 70 ml |
| 90 ml | 3 | 100 ml |
| 120 ml | 4 | 135 ml |
| 150 ml | 5 | 170 ml |
| 180 ml | 6 | 200 ml |
| 210 ml | 7 | 235 ml |

Breast milk is the best nutrition for a baby. Follow on formula should only be used upon the advice of a health care professional.

Important

Wash your hands before you start cooking the infant formula. Always prepare the mixture with fresh, boiled and chilled water and always use clean sterile bottles. Only prepare one meal at a time.

Measuring

It is important for your baby's health to follow the instructions on the pack and to use only the special scoop provided in the pack when measuring out the powder.

Do not take more or less powder.



HiPP Customer Service NO tel. +46 8 799 1926 www.hipp.no

